



Palladium Acetate

Unleashing Catalytic Potential




HERAEUS – YOUR EXPERT FOR PALLADIUM ACETATE

INTRODUCTION

Catalyzed reactions, such as coupling reactions, aminations or carbonylations, play an important part in the syntheses of sophisticated active ingredients for pharmaceutical, agro- and fine-chemistry. Many of these reactions are run with the help of homogenous palladium catalysts. Palladium acetate, $\text{Pd}_3(\text{OAc})_6$,

is one of the most common Pd sources in this type of catalysis. It is a well defined, brown compound with good solubility in typical organic solvents. It serves as a reliable source of soluble palladium, which can then interact with additional ligands in the reaction mixture.

COMMON COUPLING REACTIONS WITH PALLADIUM ACETATE



As a prime example, Palladium acetate finds extensive application in Suzuki-Miyaura cross-coupling reactions, which allow for the formation of carbon-carbon bonds between aryl or vinyl boronic acids and aryl or vinyl halides. The reaction conditions involve the use of a base, such as sodium or potassium carbonate, along with a phosphine ligand, commonly triphenylphosphine (PPh₃). Suitable solvents for Suzuki-Miyaura cross-couplings include toluene or dioxane.

Further, it is highly effective in Heck reactions, a significant carbon-carbon bond-forming reaction. When used as a catalyst, typically in a palladium acetate/PPh₃ (triphenylphosphine) system, it enables the coupling of aryl or vinyl halides with various olefins.

The reaction conditions typically involve using a base, such as a carbonate or hydroxide, and an appropriate solvent such as N,N-dimethylformamide (DMF) or tetrahydrofuran (THF).

In Sonogashira couplings, another carbon-carbon bond-forming reaction, Palladium acetate serves as an effective catalyst for the coupling of terminal alkynes with aryl or vinyl halides. It is often used in combination with a copper(I) salt as a co-catalyst, such as copper(I) iodide (CuI), facilitating the formation of the desired products. The reaction conditions typically involve the use of a base, such as potassium carbonate, and solvents such as dichloromethane (DCM) or toluene.


HERAEUS – YOUR EXPERT FOR PALLADIUM ACETATE

REACTIONS

- CC-Coupling Reactions
- CN-Coupling Reactions
- CH Activations
- Carbonylations

APPLICATIONS

- ✓ Agro chemicals
- ✓ Fine chemicals
- ✓ Pharmaceuticals



SPECIFICATIONS

Molecular formula: $\text{Pd}_3(\text{OAc})_6$

Molecular Weight: 672


CAS Number: 3375-31-3

Metal content: 46–48 %

Solubility: Acetone, methanol, acetonitrile, dichloromethane, chloroform

Description: Palladium acetate trimer

EXEMPLARY FT-IR SPECTRUM



Palladium acetate stands as a powerful catalyst, offering excellent solubility in common organic solvents, high reactivity, selectivity, and versatility. Its solubility allows for the preparation of homogeneous catalyst solutions, ensuring optimum performance and ease of handling in various organic reactions.

UNLOCK THE CATALYTIC POTENTIAL WITH PALLADIUM ACETATE!

ABOUT HERAEUS PRECIOUS METALS

Heraeus Precious Metals is a leading provider of precious metals services and products. We combine all activities related to our comprehensive expertise in the precious metals loop – from trading to precious metals products to recycling.

Heraeus Precious Metals is one of the world's largest refiners of platinum group metals (PGMs) and a leading name in industrial precious metals trading. Our precious metals products are used in a wide variety of industries,

including the chemical, pharmaceutical, glass, electronics and automotive industries. We offer top quality solutions and products based on many years of experience and technical expertise. We are a reliable development partner for our customers and find the best solutions for their requirements.

Heraeus Precious Metals

Heraeus Precious Metals GmbH & Co. KG
Heraeusstr. 12–14
63450 Hanau, Germany
chemicals@heraeus.com

www.heraeus-precious-metals.com